

Antimony Anomalies around abandoned Silver Mines in Tyrol, Austria

Christian Wolkersdorfer, Thomas Wackwitz

TU Bergakademie Freiberg
Lehrstuhl für Hydrogeologie
Gustav-Zeuner-Str. 12, 09599 Freiberg/Sachsen, Germany
e-mail-address: c.wolke@imwa.info

Abstract

Austria was one of the richest Silver producers in Medieval Times, with most of the important mines located in Tyrol. Nearly all those mines in the Inntal Valley mined Fahlores of which the Silver contents were the basis for Tyrol's political power by that time. In 1999 a large rock slide caused the last working mine to shut down, and intensive geotechnical, geological, and hydrogeological investigations started around the sliding area. These investigations found Sb anomalies in many springs around the dolomite underground mine and in the underground mine itself. While some scientists recognized a close relation to the former Silver mine, the mine owner suggested a relation to the geological situation.

About 50 sampling points in different geological strata near the Großkogel and Schwaz dolomite and visitor's mines were sampled in summer and winter and analysed for the main and trace elements. As a result of those investigations we were able to show that the anomalies are closely linked to the geological situation and that mined as well as unmined areas are subject to Sb anomalies of up to 2.2 mg L^{-1} .

It is clear from our investigations that the Antimony anomalies are linked to the geological situation and that the current operators are not responsible for the Antimony concentrations that can be found today.

1 Introduction

1.1 Locations and Mining History

One of the most important Austrian silver production regions in medieval and early modern times was Tyrol. Between Schwaz and Brixlegg, both located in the Inntal valley north-east of Innsbruck, numerous silver mines existed that mined silver-bearing fahlore (Fig.1). Both Tyrol's political power and the economical power of the Augsburg Fugger family are due to the silver mined in the region. The investigations described here concentrated on the Schwaz and the St. Gertraudi mining districts, covering an area of 10 km² and 5 km² respectively. Originally the investigations were carried out to conduct mine water tracer tests in the Georgi-Unterbau mine south-east of Brixlegg, but after identifying the antimony anomalies, they were extended to cover the Schwaz area as well.

Two main adits were used to dewater the mined areas of St. Gertraudi and Schwaz: the Georgi-Unterbau adit and the Wilhelm-Erbstollen adit. Those two adits were the basis for the underground sampling described here.

Mining in the Schwaz area dates back to Bronze Age times (Rieser and Schrat-




Fig. 1: Location of the mining and investigation areas of Schwaz (left) and Großkogel (right) in central Tyrol, Austria. Mining symbols mark the entrances to the two main dewatering adits.

tenthaler 2002) and started again in the 13th century (Hanneberg and Schuster 1994; Palme et al. 2002). By far the most important mining area was the Falkenstein mining area limited by the Bucher Bach in the east and the Lahnbach in the west. Details about the mining in the surroundings of Brixlegg can already be found in a goods list from the year 1416. Many miners from Bohemia, Saxony and other Central European mining regions came to the Inn valley, attracted by the message of rich silver findings. In 1427 Duke Friedrich released a mining order for Schwaz to regulate the pit lending and the operational procedures. The freedom of mining promoted the prosperity of the mining area. Twenty-six pits were working at the Falkenstein before the year 1460, and within the years to 1499 another 231 mining leases were awarded. Because of this development more than 10,000 miners were employed in over 1,000 galleries and Schwaz became the most important silver mining area in Europe. Thus, the Falkenstein mines achieving the highest production with a peak of 15.7 t of fine silver in 1523. Due to mismanagement and the increasing import of silver from the Americas, silver mining in the Schwaz area ended in the late 17th century.

As a result of exploration in the Schwaz area, in 1873 the Wilhelm Erbstollen was driven to undermine the Falkenstein area. Rich fahlores were found in the Krummörter pit, but ore mining ceased in 1957. Since then the Montanwerke Brixlegg AG mined high quality dolomite until a mountain collapse caused the mines closure by 1999.

In 1887 the miners started to drive the Georgi-Unterbau at St. Gertraudi, which opened up rich fahlore and barite resources. Starting in 1900, a 100 m deep two compartment blind shaft was sunk, connecting the 20, 40, 70 and 100 m main levels and the 10, 75 and 80 m sublevels (Pirkl 1961; Mutschlechner 1984; Krischker 1990; Hanneberg and Schuster 1994) as well as the 14-Nothelfer pit and the Barbara pit with each other. It was partially to fully flooded during the 1950's (Hießleitner 1951; Schmidegg 1953) and pumped out again in 1984 and 1988 for ore prospecting (Krischker 1990). Since 1990 the blind shaft has been flooded and therefore stationary hydraulic conditions exist.

1.2 Geological situation

Most parts of the mines are within the Devonian Schwaz Dolostone (*Schwazer Dolomit*) of the Northern Tyrol Greywacke area (*Nordtiroler Grauwackenzone*). Typically, the Schwaz Dolostone is a very hard, light white to light grey dolostone, being highly brecciated and fissured in the area investigated. The dolostone hosts silver and mercury bearing fahlores (there under Schwazite) as well as barite, the mineralisation being bound, but not restricted, to the breccia zones (Pirkl 1961; Wöbking 1982). Grundmann and Martinek (1994), Schnorrer (1994) and Schnorrer (1996) describe 20 ore minerals being characteristic for the Schwaz Dolostone and a total of over 132 minerals for the Schwaz-Brixlegg area, there under REE minerals. According to Arlt and Diamond (1998), who investigated fahlores of the Schwaz-Brixlegg mining area, the fahlores of the Georgi-Unterbau comprise of 41% Cu; 0.5% Ag; 2% Fe; 5% Zn; 0.7% Hg; 0.02% Mn; 16% Sb; 9%

As; and 26% S. Unfortunately, no data on REE concentrations in the rocks of the Greywacke and their interpretation can be found in the literature.

A first physico-chemical investigation of the flooded blind shaft was conducted in December 2000. In addition to temperature and salinity measurements water samples were taken to be analysed on site and in the laboratory. Further investigations, including two tracer tests with 15 µm microspheres and Na-fluorescein, were conducted in August 2001 and February 2002 (Wolkersdorfer et al. 2002; Wolkersdorfer 2002).

1.3 Eiblschrofen Rock Slide

On July 10th 1999 a large rock slide occurred at the Eiblschrofen above the city of Schwaz. Several hundred inhabitants had to be evacuated and the Schwaz dolomite mine (run by Montanwerke Brixlegg AG) as well as the Schwaz visitor mine (run by Silberbergwerk Schwaz Besucherführungsgesellschaft m.b.H.) had to be closed immediately (Weber and Schneider 2000). The visitor mine subsequently re-opened, but the dolomite mine is still closed as a result of missing authority permissions.

As a result of the rock slide intensive investigations, including the hydrogeological situation, were initiated. Those investigations showed unusually high antimony concentrations in the surface and mine waters of the Schwaz mining area (Millen 2003). It was assumed by the authorities that these anomalies were a result of the current mining operations. Therefore the dolomite mine operator initiated a hydrogeological sampling programme to investigate the natural antimony background of the Schwaz and the Großkogel mining areas.

The aim of those hydrogeological investigations was to find relationships between the lithology and antimony anomalies, the aerial distribution of the antimony concentrations and potential remediation methods for the mine water. Furthermore, it was hoped that it could be shown whether there was a potential anthropogenic cause for the anomalies.

2 Investigations

A total of 138 water samples from the Großkogel and the Schwaz mining areas were used for the investigations described here. 98 samples were collected by the authors, 21 are cited by Weber and Schneider (2000) and 19 samples were added by the Montanwerke Brixlegg AG. Details of the sampling procedure can only be given for the 98 samples collected by the authors. They were collected from 29 sampling points in the Schwaz area and 19 from the Großkogel area, there under 27 surface and 21 underground sampling points. All sampling points were chosen to represent the main rock types outcropping at the Großkogel and around Schwaz at the surface and the main underground workings in the Großkogel barite and the Schwaz dolostone mines. Furthermore, depth dependent antimony profiles were measured in the 100 m deep Georgi Unterbau shaft.

All samples were filtered on-site with a 0.45 µm cellulose acetate filter and decanted into 500 mL bottles for the main ions and 50 mL bottles for the trace elements, the latter being acidified for stabilisation. All bottles were labelled with the sample number, the date, and the sampling point and stored in a cold and cool place thereafter. On site the pH, conductivity, temperature, redox-potential, oxygen-content, acidity (base capacity k_B 8.2), alkalinity (acid capacity k_S 4.3), and the iron concentrations (ferric and ferrous) were measured. Oxygen was measured with a WTW (Wissenschaftlich Technische Werke, Weinheim/Germany) oxygen sensor, iron with a Hach DR890 field photometer (Hach Company, Loveland CO/USA), and all other parameters with an Ultrameter 6P (MYRON L Company, Karlsbad/USA).

Main ions (Li, Na, K, Ca, Mg, Cl, NO₃, SO₄) were measured by ion chromatography with a D6000 of Merck-HITACHI, trace elements (Ag, As, Au, Ba, Bi, Cd, Co, Cs, Cu, Hg, Mn, Mo, Ni, Pb, Pd, Rb, Sb, Sc, Se, Sn, Sr, U, W, Zn) with the ICP-MS of the Fakultät Forst, Geo- und Hydrowissenschaften (Tharandt; Dresden University). Fluoride was measured with ion selective probes F500 (WTW Weinheim) and reference electrode SE 20/EB (Sensortechnik Meinsberg) as well as probe 94-09 (Orion, Boston, USA; with reference electrode included).

To compare the reliability of the low antimony concentrations in the river Inn, the relevant samples were also measured by neutron activation analyses (Freiberg Institute for Archeometry, Prof. E. Pernicka).

Using these results, statistical (SPSS) and chemico-thermodynamical (PHREEQC) investigations were carried out to interpret the data obtained. In addition an extensive literature review was conducted, which showed that antimony anomalies in mine and surface water had not previously been of concern.

3 Results

European and Austrian regulations require an antimony content of 5 µg L⁻¹ in drinking water (EU 98/83; TWV BGBl 304/2001). No regulations exist for surface waters, but the German Länderarbeitsgemeinschaft Wasser (LAWA) published a list of antimony values which require inspection or action when exceeded in groundwaters (Länderarbeitsgemeinschaft Wasser [LAWA] 1994). Thereafter, inspections are necessary if the antimony concentrations exceed 2–10 µg L⁻¹ and remediation actions are required if 20–60 µg L⁻¹ are exceeded.

According to investigations conducted by the Austrian Environment Agency (Philippitsch and Grath 2003), antimony in Austrian karst waters has a mean of 0.2 µg L⁻¹ and a maximum of 0.7 µg L⁻¹ ($n = 50$). Drinking waters in Austria are regularly below the drinking water standards. This complies with other investigations, which have shown that surface water concentrations range between < 0.05 and 0.08 µg L⁻¹, ground water between < 0.02 and 1.33 µg L⁻¹ and mine waters between < 0.1 and 250 µg L⁻¹ (Mattheß 1994; Merkel and Sperling 1998). Extreme antimony concentrations were observed in brines or thermal waters, reaching 8–1,000 µg L⁻¹ (Wedepohl et al. 1969–1978).




Fig. 2: Hydrogeochemical classification of the analysed surface and mine waters in the PIPER-diagram (explanation in the text).

Both the Schwaz and the Großkogel areas revealed mean and maximum antimony concentrations which are far higher than those reported above (Table 1). Though geologically slightly different, the means and maxima for both areas are very similar (539 and $512 \mu\text{g L}^{-1}$; 1768 and $1758 \mu\text{g L}^{-1}$, respectively). Other sources also report antimony concentrations of up to $2200 \mu\text{g L}^{-1}$, but no particular information about the sampling procedure is given, therefore those values will not be included in the detailed interpretation (Montanwerke Brixlegg AG; Weber and Schneider 2000).

Hydrogeochemically the surface and ground waters of Schwaz and Kogel are very similar, plotting into the Ca-Mg-HCO₃ and the Mg-Ca-HCO₃ fields (Fig. 2). Only some of the Schwaz mine waters are enriched in sulphate and therefore plot into the Ca-Mg-HCO₃-SO₄ and Mg-Ca-HCO₃-SO₄ fields. This can also be seen from Table 1, with the mean Schwaz sulphate concentrations being twice as high as the Kogel values. With principal component analyses the relationship of all parameters analysed were investigated. It could clearly be seen that Cu, Sb, As, and Zn, the main metals in the fahlores, plot into the same field. Ag, Bi, Mn, Pb, Cd, and Hg, and some of the other metals in the fahlore, plot in different areas, because their hydrogeochemical behaviour differs from their behaviour during rock and mineral formation processes. Their relative concentrations in the waters equal those of the fahlores (Fig. 3).

Antimony Anomalies around abandoned Silver Mines in Tyrol, Austria




Fig. 3: Box plots of the main heavy and semi-metal concentrations selected in accordance with their abundance in the Schwaz and Kogel fahlores. All concentrations in mmol L⁻¹; the small numbers under the elements show the concentration in µg L⁻¹.




Fig. 4: Box plots of antimony in relation to the five main lithological units of Schwaz. TWV: standard of Austrian drinking water regulation; LAWA: standard for action.

46 water samples of the Schwaz area could directly be assigned to one of the five rock types outcropping there: Schwaz Dolomite (8 samples); Triassic Buntsandstein (6 samples); Phyllites (18 samples); Triassic Limestones (4 samples) and Quaternary Sediments (2 samples). Only concentrations of the samples from the Quaternary Sediments fell below the Austrian drinking waters standards, while the

samples from the phyllites and the Triassic Limestones were under the German LAWA limits. More than half of the samples taken from the Schwaz Dolomite and the Buntsandstein exceeded the LAWA standard (Fig. 4).

In the mine waters of the Schwaz dolomite mine antimony concentrations above $600 \mu\text{g L}^{-1}$ were related to the old workings in the “Abbau I – III”, and in the Georgi Unterbau to the flooded shaft. All the other mine waters were below $600 \mu\text{g L}^{-1}$, and in the Georgi Unterbau below $200 \mu\text{g L}^{-1}$. At the dewatering adits of the Wilhelm Erbstollen and the visitor’s mine the antimony concentrations were $326 \mu\text{g L}^{-1}$ and $185 \mu\text{g L}^{-1}$ respectively. More important than the concentrations are the mass loads. At “Abbau III”, with the highest antimony concentration, a load of 0.4 kg day^{-1} was recorded, and at the dewatering adit of the old workings of the “Messerschmitt Halle” a load of approximately 13 kg day^{-1} was measured. For comparison, the Sb loads of the 3 main adits are 0.2 kg day^{-1} for the visitor’s mine, 16 kg day^{-1} for the Wilhelm Erbstollen, and 0.1 kg day^{-1} for the Georgi Unterbau.

4 Discussion

It has been shown that the high antimony anomalies are closely linked to the Schwaz Dolomite, which is the host rock for the silver bearing fahlores. To a lesser extent, higher antimony concentrations are associated with the Buntsandstein outcrops at the surface. All the other lithologies investigated may contain local springs with antimony contents exceeding the LAWA standards, although in some cases it was not clear whether the springs might actually arise from abandoned adits now covered by rock debris. Antimony concentrations in waters from the Triassic limestones and the Quaternary Sediments were regularly below the limits of the drinking water regulations.

Within the mine workings, both in the Schwaz and the Kogel cases, high antimony contents are clearly linked to the abandoned mine workings and to the “Abbau III”, which was subject to a surface collapse in 1995. In this area, surface water flows through old abandoned mine workings of medieval and Bronze Age times and is therefore enriched in antimony and in sulphate. The new workings in “Abbau IV” had antimony concentrations lower than $200 \mu\text{g L}^{-1}$ and the extreme mine water inflows in the Georgi Unterbau away from the flooded shaft also had low antimony concentrations.

It can therefore be concluded that the high antimony concentrations are clearly linked to the Schwaz Dolomite and to the abandoned mine workings which were not the subject of dolomite mining of the Montanwerke Brixlegg AG. Nonetheless, in case of a continuation of the dolomite mining, the operator shall ensure that the contact time between the fahlore bearing dolomite and the mine water is as short as possible. By using flexible pipes at the most dominant water inflows, and foils to protect the excavated material from “drippers”, the antimony load might be reduced significantly. In the case of the three dewatering adits it is advisable to use the area owned by the operators for constructing wetlands which may

Antimony Anomalies around abandoned Silver Mines in Tyrol, Austria

parameter	Kogel			Schwaz			<i>n</i>
	min	max	Mean	min	max	mean	
Temperature; °C	7.5	17.2	9.2	3.4	15.2	10.5	94
pH; –	6.9	8.4	7.8	7.3	8.9	8.0	100
Conductivity; $\mu\text{S}/\text{cm}$	230	486	393	66	1107	407	101
Redox, mV	320	530	450	270	480	440	80
HCO ₃ ; mg/L	131	262	213	46	272	188	60
SO ₄ ²⁻ ; mg/L	2	45	29	3	470	67	62
Sb; mg/L	0.001	1.768	0.593	0.0002	1.758	0.512	77

Tab. 1: Main results of the hydrogeological investigations at the Kogel and Schwaz mining areas. *n*: number of measurements (varies due to sampling conditions).

serve to reduce metal concentrations. Taking into account that arsenic and antimony have a similar chemical behaviour this passive treatment technique for antimony may be an effective remediation measure.

Acknowledgements

We would like to thank the Montanwerke Brixlegg AG, especially Robert Stibich and Eduard Ludescher, for their support and for giving the permission to publish our results. Thanks to “Team 2000” who provided us many interesting information about the mine layout. The following people conducted analyses or helped us to clarify open questions: Ernst Pernička, Peter Gstrein, Leopold Weber, Bernhard Millen, and Johannes Bauer. Last but not least thanks to Katy Unger-Shayesteh and Andrea Hasche. Part of the work was financed by the European Commission R&D project PIRAMID (EVK1-CT-1999-000021).

References

- Arlt T, Diamond L W. 1998. Composition of tetrahedrite-tennantite and “schwazite” in the Schwaz silver mines, North Tyrol, Austria. *Mineralogical Magazine*; 62(6):801–820.
- Grundmann G, Martinek K-P. 1994. Erzminerale und Gangarten des Bergbauebietes Schwaz-Brixlegg. *Mineralien Magazin, Lapis*; 19(7–8):28–40.
- Hanneberg A, Schuster H. 1994. Geschichte des Bergbaus in Schwaz und Brixlegg. *Mineralien Magazin, Lapis*; 19(7–8):13–21.
- Hiebleitner G. 1951. Geologischer Gutachtensbericht über den Schwespatbergbau Großkogel bei Brixlegg. Unveröffentlichtes Gutachten; Archiv Montanwerke Brixlegg.
- Krischker A G. 1990. Die Baryt-Fahlerz-Lagerstätte St. Gertraudi/Brixlegg. Innsbruck: Unveröff. Dipl.-Arb. Univ. Innsbruck: 206.
- Länderarbeitsgemeinschaft Wasser (LAWA). 1994. Empfehlungen für die Erkundung, Bewertung und Behandlung von Grundwasserschäden. Stuttgart: Länderarbeitsgemeinschaft Wasser: 19.

- Mattheß G. 1994. Die Beschaffenheit des Grundwassers. In: Mattheß G.. Lehrbuch der Hydrogeologie, vol. 2, 3 edn. Berlin u.a.: Borntraeger: 499.
- Merkel B, Sperling B. 1998. DVWK Schriften – Hydrogeochemische Stoffsysteme Teil 2. In: Deutscher Verband für Wasserwirtschaft und Kulturbau e.V. (DVWK), vol. 117. Bonn: Kommissionsvertrieb Wirtschafts- und Verlagsgesellschaft Gas und Wasser mbH: 1–415.
- Millen B M J. 2003. Aspects of the Hydrogeology of a Mining Region with a Focus on the Antimony Content of the Spring-water, Eiblschrofen Massif, Schwaz, Tyrol, Austria. Mitteilungen der Österreichischen Geologischen Gesellschaft; 94:139–156.
- Mutschlechner G. 1984. Erzbergbau und Bergwesen im Berggericht Rattenberg. Innsbruck: Eigenverlag Gemeinden Alpbach, Brixlegg, Rattenberg und Reith im Alpbachtal: 163.
- Palme R, Gstrein P, Ingenhaeff W. 2002. Glück Auf! – Faszination Schwazer Silberbergwerk. Innsbruck: Berenkamp: 111.
- Philippitsch R, Grath J. 2003. Wassergüte in Österreich Jahresbericht 2002 - Erhebung der Wassergüte gemäß Hydrographiegesetz (BGBl.-Nr. 252/90, i.d.g.F.). Wien: Umweltbundesamt: 178.
- Pirkl H. 1961. Geologie des Trias-Streifens und des Schwazer Dolomits südlich des Inn zwischen Schwaz und Wörgl (Tirol). Jb Geol B-A; 104(1):1–150.
- Rieser B, Schrattenthaler H. 2002. Prähistorischer Bergbau im Raum Schwaz-Brixlegg. St. Gertraudi: Edition Tirol: 125.
- Schmidegg O. 1953. Die Erzlagerstätten am Reiter Kopf und am Reiter Kogel. Schlern-Schriften; 101:17–25.
- Schnorrer G. 1994. Die Sekundärminerale des Bergbaugebietes Schwaz-Brixlegg in Tirol. Mineralien-Magazin, Lapis; 19(7–8):41–68.
- Schnorrer G. 1996. Neue Sekundärminerale des alten Bergreviers Schwaz-Brixlegg in Tirol. Mineralien-Magazin, Lapis; 21(12):16–19.
- Weber L, Schneider J F. 2000. Geologie, Geotechnik, Georisiken – Internationales Expertengutachten zum Verfahren des Bundesministeriums für Wirtschaft und Arbeit nach § 179 Mineralrohstoffgesetz betreffend den Bergbaubetrieb „Dolomitbergbau Schwaz“ der Montanwerke Brixlegg Aktiengesellschaft und das Schaubergwerk „Silberbergwerk Schwaz“ der Silberbergwerk Schwaz Besucherführungsgesellschaft m.b.H. in Zusammenhang mit den Felssturzereignissen am Eiblschrofen. In: Österreichisches Bundesministerium für Wirtschaft und Arbeit. Wien: Bundesministerium für Wirtschaft und Arbeit, Universität für Bodenkultur: 107.
- Wedepohl K H, Correns C W, Shaw D M, Turekian K K, Zemann J. 1969–1978. Handbook of Geochemistry. Berlin Heidelberg New York: Springer.
- Wöbking H. 1982. Untertägige Eigenpotentialmessungen in den Bergbauen Grosskogel bei Brixlegg und Falkenstein bei Schwaz. Berg- und Hüttenm Mh; 127(12):476–483.
- Wolkersdorfer C. 2002. Rare Earth Elements (REEs) as Natural Tracers in Mine Waters. In: Merkel B., Planer-Friedrich B., Wolkersdorfer Ch. Uranium in the Aquatic Environment. Heidelberg: Springer: 967–974.
- Wolkersdorfer C, Hasche A, Unger K, Wackwitz T. 2002. Tracer Techniken im Bergbau – Georgi-Unterbau bei Brixlegg/Tirol. Wissenschaftliche Mitteilungen; 19:37–43.